

Many-body systems in the presence of the random interaction and the J pairing
interaction

Akito Arima

The House of Councilors, 2-1-1 Nagatacho, Chiyodaku, Tokyo 100-8962, Japan

In this talk I shall discuss some regularities of many-body systems in the presence of random interactions and regularities of a single- j shell for the J pairing interaction which works only when two particles are coupled to spin J . I shall first explain an empirical rule to predict the spin I ground state probability. Then I shall present some interesting results of a single- j shell under the J pairing interaction. Last I shall discuss some preliminary results of binding energies in the presence of random two-body interactions.

PACS number: 21.60.Ev, 21.60.Fw, 24.60.Lz, 05.45.-a

It is my great pleasure for me to talk to you here. I would like to thank the organizers of this conference and say “congratulations” to Professor Stuart Pittel. I am also very glad to see many of my friends.

My talk includes three subjects.

- (1) spin 0 ground state dominance;
- (2) Some regularities under the J pairing interaction;
- (3) ground state parities and binding energies.

1 spin 0 ground state dominance

The spin 0^+ ground state (0 g.s.) dominance was discovered and first studied by Johnson, Bertsch, Dean, and Talmi [1]. This phenomenon has attracted much attention [2].

In Ref. [3] we proposed a simple approach to predict the probability (denoted as $P(I)$) for a certain spin I to be the ground state spin without using random interactions: We first set one of the two-body matrix elements of the problem to -1 and all the rest to zero, and then see which angular momentum I gives the lowest eigenvalue among all eigenvalues of this many-body system. If the number of independent two-body interaction matrix elements is N , the above procedure is repeated N times. After all N calculations have been done, we count how many times (denoted as $\mathcal{N}_I$) the angular momentum I gives the lowest eigenvalue. We then predict $P(I)$ to be $\mathcal{N}_I/N$.

This approach was applied to predict the $P(I)$ for various systems, such as fermions in a single- j shell or many- j shells, sd bosons or sdg bosons. Here we take a single- j shell with $j = 9/2$ as an example. The Hamiltonian is defined as follows.

$$H = \sum_J G_J A^{J\dagger} \cdot A^J \equiv \sum_J \sqrt{2J+1} \left(A^{J\dagger} \times A^J \right)^{(0)},$$

$$A^{J\dagger} = \frac{1}{\sqrt{2}} \left[a_j^\dagger \times a_j^\dagger \right]^{(J)}, \quad A^J = -(-1)^M \frac{1}{\sqrt{2}} \left[\tilde{a}_j \times \tilde{a}_j \right]^{(J)}. \quad (1)$$

where G_J is given by $\langle j^2 J | V | j^2 J \rangle$, and V is a two-body interaction. Here the J

pairing interaction is defined as the interaction which has $G_{J'} = -\delta_{JJ'}$.

One can obtain that $\mathcal{N}_0=3$ and $\mathcal{N}_4=\mathcal{N}_{I_{\max}}=1$ for four fermions ($n = 4$, n labels the particle number) and $\mathcal{N}_j=2$ and $\mathcal{N}_{3/2}=\mathcal{N}_{5/2}=\mathcal{N}_{I_{\max}}=1$ for $n = 5$. Here $N = j + 1/2 = 5$. Thus one predicts that $P(0)=60\%$ and $P(4) = P(I_{\max}) = 20\%$ for $n = 4$, and $P(j)=40\%$ and $P(3/2) = P(5/2) = P(I_{\max}) = 20\%$ for $n = 5$. The I g.s. probabilities obtained by using the random interactions are as follows: $P(0)=66.4\%$, $P(4) = 11.8\%$, $P(I_{\max}) = 17.9\%$ and other $P(I)$'s are close to zero for $n = 4$; $P(j)=33.9\%$ and $P(3/2) = 20.5\%$, $P(5/2) = 15.5\%$, $P(I_{\max}) = 18.4\%$ and other $P(I)$'s are close to zero for $n = 5$. The agreement between the predicted values and those obtained by the random interactions is good.

There are two other conclusions based on our empirical rule. First, $P(I_{\max})$ is considerably large for a single- j shell, with the predicted $P(I_{\max}) = 1/N = 1/(j + \frac{1}{2})$ which is independent of n . This prediction has been confirmed quantitatively. Second, through the process of finding the spin I of the ground state of the system with $G_{J'} = -\delta_{JJ'}$, one is able to predict which interactions are essential for a spin I g.s. probability.

2 Regularities of many-body systems under the J pairing interaction

The above empirical rule stimulated our studies of a single- j shell under the J pairing interaction. To avoid confusion, here we use superscript (n) to specify the particle number n for spin $I^{(n)}$ and the eigenvalue $E_{I^{(n)}, J(j)}^{(n)}$ for a single- j shell with the J pairing interaction.

We recently showed in Ref. [4] that a system of three fermions in a single- j shell is exactly solvable for any J pairing interaction. There is at most one non-zero eigenvalue for a fixed J and for any $I^{(3)}$. The non-zero eigenvalue is given by

$$E_{I^{(3)}, J(j)}^{(3)} = -1 - 2(2j + 1) \left\{ \begin{matrix} J & j & I^{(3)} \\ J & j & j \end{matrix} \right\}. \quad (2)$$

for $G_{J'} = -\delta_{JJ'}$.

The summation of all eigenvalues over J for a fixed $I^{(3)}$ is equal to the number of spin $I^{(3)}$ states multiplied by a factor $\frac{n(n-1)}{2} = 3$. Combining this with the result of Eq. (2), one can obtain a number of new sum rules for six- j symbols. These sum rules were found and discussed in Ref. [4].

The case of four fermions for the J pairing interaction is not soluble except for $J = 0$. Here we concentrate on the $J_{\max}$ pairing. In Ref. [5] we found that many of eigenvalues $E_{I^{(4)}, J_{\max}(j)}^{(4)}$ are asymptotically integers (0, -1 and -2), and related them with the number of pairs with spin $J_{\max}$ in their wavefunctions. Besides these asymptotic integer eigenvalues, there are many $E_{I^{(4)}, J_{\max}(j)}^{(4)}$ which are not close to integers. We called them “non-integer” eigenvalues, and noticed that they are very close to the non-integer eigenvalues of $n = 3$. Therefore, it might be anticipated the wavefunction corresponding to the non-integer $E_{I^{(4)}, J_{\max}(j)}^{(4)}$ could be approximated by that of the corresponding $E_{I^{(3)}, J_{\max}(j)}^{(3)}$, to which $E_{I^{(4)}, J_{\max}(j)}^{(4)}$ is asymptotically equal, coupled to a single- j particle. This anticipation was confirmed by calculating their overlaps which are 1.0 within a high precision. This last particle is called a “spectator” because of its weak coupling with the other strongly bound three particles which can be regarded as a cluster.

It is noted that the similar pattern holds for $n = 5$ and 6, and also for bosons with spin l . The non-integer eigenvalues and integer eigenvalues can be unified by the following picture of clusters which include both $J_{\max}$ pairs and spectators: The $J_{\max}$ pairing interaction favors clusters with n_1 particles ($n_1 \leq n$), with $I^{(n_1)} \sim I_{\max}^{(n_1)}$.

As is well known, the existence of degeneracy indicates that the Hamiltonian has a certain symmetry. The degeneracy for $J_{\max}$ pairing interaction, however, is not exact. It would be interesting to explore the broken symmetry hidden in the $J_{\max}$ pairing interaction discussed in this paper. It would be also interesting to discuss the modification of the $J_{\max}$ pairing interaction in order to recover the exact degeneracy.

3 Ground state parities and binding energies

There are many interesting patterns of many-body systems in the presence of random interactions. Some characteristic features such as the spin 0 g.s. dominance and the odd-even staggering of binding energies have been numerically well known under the random interactions. Below I discuss two other regular patterns.

The first is related to the parity distribution. In Ref. [6] we studied many systems filling different shells in the presence of two-body random ensemble, and found that positive parity is dominant in the ground states of even-even nuclei, and the possibility for the ground states of odd-mass and odd-odd nuclei to have either positive or negative parity is comparable in general. This feature is very similar to that for the 0 g.s. dominance. Because parity is a much simpler quantity, the positive parity dominance might be well understood in the near future. This may help us to obtain a deeper understanding of the 0 g.s. dominance.

The second is related to binding energies obtained by the random interactions. The binding energy is a very important and fundamental quantity not only for various aspects of nuclear physics but also for other branches of physics such as astrophysics. For a recent review of nuclear masses, see Ref. [7]. Recently Bohigas and Leboeuf claimed in Ref. [8] that there is a lower limit on the accuracy of theoretical predictions of binding energies due to chaotic components in the nuclear mass. Here I present some preliminary results for binding energies of a few neighboring nuclides in the *sd* shell which are obtained by using the random interactions. We are interested in the Garvey-Kelson relations which connect the masses of neighboring nuclides. Our calculated results seem to satisfy the algebraic Garvey-Kelson relations within a reasonable precision in the presence of random interactions. For instance, we calculated the deviations from the Garvey-Kelson relation for example around ^{22}Na , which involves of binding energies of $\text{Na}^{21,22}$, $\text{Mg}^{21,23}$ and $\text{Al}^{22,23}$, and found that these deviations are small. More work is necessary to clarify the indications of Ref. [8].

4 Summary

In this talk I have discussed a few interesting aspects concerning the regularities of many-body systems under the random interactions and a single- j shell under the J pairing interaction. First, I explained our empirical rule which is useful in predicting the I g.s. probabilities and understanding the spin 0 g.s. dominance. Second, I showed that three fermions in a single- j shell for any J pairing interaction is solvable. I also discussed more fermions among which the $J_{\max}$ pairing interaction works, and suggested that the $J_{\max}$ pairing interaction favors clusters, where $I^{(n_1)}$ ($n_1 \leq n$) for each cluster is very close to or equal to the maximum spin for n_1 particles. Last, I showed two other interesting features for the ground state parities and binding energies by using the random interactions: (1) the positive parity dominates in the ground states of even-even nuclei, and (2) the calculated binding energies satisfy the Garvey-Kelson relations reasonably well.

Acknowledgement: I would like to thank Dr. Y. M. Zhao for his collaboration in this work. I also thank Dr. I. Talmi for his valuable comments on some results of Sec. 2.

References

- [1] C. W. Johnson, G. F. Bertsch, D. J. Dean, Phys. Rev. Lett. **80**, 2749(1998);
C. W. Johnson, G. F. Bertsch, D. J. Dean, and I. Talmi, Phys. Rev. **C61**,
014311(1999).
- [2] V. Zelevinsky and A. Volya, Phys. Rep. **391**, 311 (2004); Y.M. Zhao, A. Arima,
and N. Yoshinaga, nucl-th/0311050, submitted to Phys. Rep.
- [3] Y. M. Zhao, A. Arima, and N. Yoshinaga, Phys. Rev. **C66**, 034302 (2002).
- [4] Y. M. Zhao and A. Arima, nucl-th/0404041.
- [5] Y. M. Zhao, A. Arima, J. N. Ginocchio, and N. Yoshinaga, **C68**, 044320 (2003).

- [6] Y. M. Zhao, A. Arima, N. Shimizu, K. Ogawa, N. Yoshinaga, and O. Scholten, nucl-th/0404042, submitted to Phys. Rev. **C**.
- [7] D. Lunney, J. M. Pearson, and C. Thibault, Rev. Mod. Phys. **75**, 1021 (2002).
- [8] O. Bohigas and P. Leboeuf, Phys. Rev. Lett. **88**, 092502 (2002).